

On modified symmetries for the HIV AIDS epidemic model equations

A.W.H Mochaki¹ & J.M Manale²

¹Department of Mathematics and Applied Mathematics, University of Limpopo

Private BagX1106,Sovenga , 0727, Limpopo Province Republic of South Africa
Andrew.Mochaki@ul.ac.za

²Department of Mathematical Sciences, University of South Africa

Florida,1709, Johannesburg, Gauteng Province
Republic of South Africa
manaljm@unisa.ac.za

Keywords: HIV Aids model, Lie symmetry, Invariant solutions, Modified symmetries

In this contribution we apply Lie group theoretical methods to a system of first-order ordinary differential equations modelling the HIV/AIDS epidemic [1],[2]. In particular, we determine modified local one-parameter point symmetries, i.e., the smart symmetries, and deduce corresponding group invariant solutions.

References

- [1] Anderson RM, The role of mathematical models in the study of HIV transmission and the epidemiology of AIDS, Journal of Acquired Immune Deficiency Syndrome 1(3),1988, 241-265
- [2] Torrisi V and Nucci MC, Application of the Lie group analysis to a mathematical model which describe HIV transmission: The geometrical study of differential equations, Contemporary Mathematics, American Mathematical Society 285,2001, 11-20